

Dietary intake, physical activity and nutritional status in adults: the French nutrition and health survey (ENNS, 2006–2007)

Katia Castetbon^{1,2*}, Michel Vernay¹, Aurélie Malon¹, Benoit Salanave¹, Valérie Deschamps¹, Candice Roudier¹, Amivi Oleko¹, Emmanuelle Szego¹ and Serge Hercberg^{1,2,3}

¹Unité de Surveillance et d'Epidémiologie nutritionnelle, Institut de Veille sanitaire, Université de Paris 13, Conservatoire national des Arts et Métiers, Bobigny, France

²Unité de Recherche en Epidémiologie nutritionnelle, U557 Inserm; U1125 Inra; Cnam; UP13, CRNH IDF, Bobigny, France

³Département de Santé publique, Hôpital Avicenne, AP-HP, Bobigny, France

(Received 10 October 2008 – Revised 20 January 2009 – Accepted 20 January 2009 – First published online 2 March 2009)

The French National Programme on Nutrition and Health (Programme national nutrition santé (PNNS)), the aim of which is to reduce nutrition-related chronic diseases, necessitates monitoring of nutritional characteristics. Our objective was to describe dietary intake, physical activity and nutritional status in a national sample of adults, especially according to current French recommendations. The study is based on a cross-sectional population-based survey using a multistage sampling design (Etude nationale nutrition santé (ENNS)). Between February 2006 and March 2007, 3115 18–74-year-old adults were included (participation rate 59.7%). Energy, macronutrient and food consumption were estimated through three randomly distributed 24 h recalls, and compared to PNNS recommendations; physical activity was described using International Physical Activity Questionnaire guidelines; anthropometry, blood pressure and biochemical measurements were assessed according to national and international references. When compared to current recommendations, intake of carbohydrates (>50% energy intake without alcohol: 26.4%), SFA (<35% total lipids: 18.5%) and total fibre (>25 g/d: 13.7%) was frequently unsatisfactory. While overall consumption of 'meat, seafood and eggs' was satisfactory, that of fruits and vegetables (≥ 400 g/d: 43.8%) and seafood (two or more servings per week: 29.9%) was frequently too low. The physical activity level was satisfactory at 63.2%. Overweight was observed in 49.3% of adults, while 30.9% were hypertensive and 44.1% had dyslipidaemia. Vitamin and iron-poor status was found to affect less than 10% of the population. Based on the ENNS survey, overall nutrition remains a problem in France. Comparison of these data with those of other countries could contribute to a better understanding of variations in nutrition-related diseases.

Nutritional surveys: Diet assessment: Physical activity: Biochemical markers: Adults

Satisfactory diet, physical activity and nutritional status are recognized as major determinants of health. Many epidemiological studies have shown a significant association between nutrition and CVD, certain cancers, type 2 diabetes and other age-related diseases⁽¹⁾. Healthcare expenses that could be spared through nutritional prevention, especially of obesity, are subject to debate⁽²⁾. Indeed, when obesity-related mortality is prevented and life expectancy increases, the cost of less lethal illnesses may also increase⁽³⁾. Nonetheless, prevention of diseases associated with nutrition remains a priority. Benefits in terms of health and life expectancy that could be expected from population intervention, *via* promotion of satisfactory nutrition, have been internationally published⁽¹⁾ and public health programmes have been widely implemented especially in Europe⁽⁴⁾.

In France, in 2002, obesity-related costs were estimated at 2.1–6.2 billion euros, i.e. 1.5–4.6% of total health expenditures⁽⁵⁾. In 2005, over 155 000 people died from cancer and

149 000 from CVD (29.5 and 28.4% of total causes of mortality, respectively)⁽⁶⁾. France appears to be relatively well protected from cardiovascular mortality compared to other European countries⁽⁷⁾, possibly due to lifestyle and environmental factors such as smoking, treatment and care, in addition to nutritional characteristics. In the mid-2000s, subjects being treated for diabetes were estimated at 3.7%⁽⁸⁾ and those treated for hypertension at 22.8%⁽⁹⁾. In 2001, the Ministry of Health implemented a national public health programme on nutrition, the 'Programme national nutrition santé' (PNNS), in order to improve population health status⁽¹⁰⁾. Public health objectives have been defined concerning diet, nutritional status and physical activity. To attain these objectives, various recommendations have been provided to the population through simple and accessible guidelines.

National surveys describing diet, physical activity and nutritional status are necessary in order to monitor changes directly or indirectly associated with public health projects.

Abbreviations: EI, energy intake without alcohol; ENNS, Etude nationale nutrition santé; IPAQ, International Physical Activity Questionnaire; PNNS, Programme national nutrition santé.

* **Corresponding author:** Dr K. Castetbon, fax +33 1 48387374, email katia.castetbon@univ-paris13.fr

In the USA, the National Health and Nutrition Examination Survey has been regularly conducted for more than three decades. Numerous analyses of these data have provided an overview of nutritional changes and the recent situation in the USA^(11,12). In Europe, despite specificities regarding nutritional behaviour, obesity and chronic disease frequencies⁽¹³⁾, such national descriptions are less systematically disseminated as a whole^(14–16). In France, data available for nutritional surveillance were disparate up until the year 2000. They separately concerned diet⁽¹⁷⁾ or body mass⁽¹⁸⁾, involved research cohorts⁽¹⁹⁾ or were limited to certain regions⁽²⁰⁾. In the framework of the PNNS, a national monitoring system dedicated to surveillance of the nutritional status of the population was specifically implemented. One of the challenges for surveillance is to obtain representative and comprehensive data that could contribute to public health policies.

In 2006–7, the French nutrition and health survey (Etude nationale nutrition santé (ENNS)) was conducted in order to describe dietary intake, physical activity and nutritional status in a national sample of adults and children living in France. The objective of the present analyses is to describe these nutritional issues in adults, especially with respect to current French recommendations.

Methods

Study design and recruitment

The ENNS survey was a cross-sectional study with a multi-stage sampling design. Recruitment was carried out throughout the entire territory of continental France for one year (February 2006 to March 2007) so as to account for seasonality of diet. The first stage of sampling involved random selection of 190 geographic zones (municipalities or groups of municipalities). This was stratified on eight large regions and on the degree of urbanization (four groups, from 'rural' to 'towns of more than 100 000 inhabitants'). The number of individuals included in each stratum was defined proportionally to the population size, with a minimum of twenty-four dwellings to be investigated. The second stage involved random selection of dwellings based on randomly generated phone lists; dwellings owned landlines, with or without phone number spread on free lists, or mobile phones only. When the detailed postal address was retrieved, an information letter was sent prior to the first phone contact with the selected household. The third stage of sampling involved selection of eligible subjects to be included in the study.

Criteria for adult eligibility were the following: residence in continental France during the period of the study, age 18–74 years, normal living accommodation for at least 5 d a week (not in an institution) with phone contact, an understanding of the issues of the study, no artificial feeding, and agreement to participate in the study. The aim of the first home visit with a study dietitian was to explain in detail the protocol of participation and to obtain agreement for participating. No incentive was given, but each subject was offered to receive his/her individual results as well as general conclusions of the study. Three runs of inclusion were distributed during the year of the study; during each run, all 190 geographical zones were investigated.

The survey protocol received the approval of the Ethical Committee (Hôpital Cochin, Paris, no. 2264), the Consultative Committee on Information Treatment of the Ministry of Research and the French Data Protection Authority (Commission nationale de l'informatique et des libertés, authorization no. 905481). Adults who participated in clinical and biochemical examinations signed informed consents separately for the scheduled analysis and for the biobank.

Data collection

Data comprised dietary intake description, clinical and biochemical marker measurements, physical activity and complementary items by questionnaires. They were collected either at home (questionnaires and, if subjects did not want to go to a health centre, anthropometry and blood pressure measurements and biochemical samples with a nurse), by phone (24 h diet recalls) or at a health examination centre of the National Health Insurance System (Caisse nationale d'assurance maladie des travailleurs salariés).

Repeated 24 h dietary recalls. Dietary intake was assessed by three 24 h recalls, one of which was on the weekend, randomly distributed within a period of 2 weeks. Trained dietitians asked subjects to describe all foods and beverages consumed the day before the interview, from midnight to midnight. Subjects gave information on food composition, recipes (especially when they were homemade) and trademark. Amounts consumed were described using a validated photography manual⁽²¹⁾ or according to standard measurements (home containers, grams indicated on the package, etc.). Data were entered on software developed for the study that included more than 3400 foods and beverages (NutriXpert[®]; Medical Expert System, Paris, France). Nutritional values for energy, macronutrients and main micronutrients came from a nutrient database previously published⁽²²⁾ and completed for recent market foods and recipes. Individuals were not previously informed of the call day so as to limit changes in diet. Information on home-added fat and addition of salt during food preparation and consumption was separately collected.

Physical activity and sedentary lifestyle. The short-form of the International Physical Activity Questionnaire (IPAQ)⁽²³⁾ in the French language was used. In brief, the numbers of days over the past week in which the subject spent more than 10 min walking and carrying out physical activity of moderate or vigorous intensity were collected. The respective durations on one of these days were collected. Physical activity description was completed by data on time spent sitting down and time spent in front of a screen (television, computer and video-games) during the past 7 d.

Anthropometry and blood pressure measurements. After the diet collection period, adults were offered the possibility of undergoing a clinical and biochemical examination. Anthropometry was measured using identical devices and according to standardized procedures adapted from WHO recommendations⁽²⁴⁾. Weight was measured to the nearest 100 g on an electronic scale (SECA[®]) with the subject lightly dressed (wearing mainly underwear) without shoes. Height was measured without shoes to the nearest 0.5 cm with a microgauge (SECA[®]) attached to the wall. Waist and hip circumferences were measured with a flexible double-graduated tape measure to the nearest 0.1 cm, and the operator was

seated at the side. Waist circumference was taken midway between the inferior margin of the last rib and the crest of the ilium.

On the same day, three measurements of diastolic and systolic blood pressure were taken on the left arm using an automatic validated device (OMRON M5-I®) and size cuff for obese subjects, when appropriate. The rest period in the same position prior to measurement was at least 5 min, and measurements were 1 min apart. The arm was placed on a support in that the elbow was in the plane of the heart. A second run of three measurements was carried out if a difference of 10 mm Hg was observed between two recordings.

Biochemical measurements. A peripheral blood sample was taken after a recommended minimum of 12 h of fasting. Eventual food and beverage consumption was collected if taken during the last 12 h. Subjects were asked to collect morning urine and store it at +4°C. Samples were then carried up to the field laboratories within 4 h after collection in frozen containers. Blood samples were immediately centrifuged when appropriate and analysed for cholesterol (total and HDL-cholesterol), TAG, glucose, Hb and ferritin. Most biochemical analyses were performed on an automated Olympus®. Serum cholesterol (total and HDL-cholesterol) and TAG were determined by enzymatic colorimetric methods; LDL-cholesterol was calculated using the *ad hoc* formula when TAG were <4.6 mmol/l⁽²⁵⁾. Plasma glucose was determined using the hexokinase method and colorimetric measurement. ELISA measurement was used for ferritin determination. Full blood count, including Hb measurement, was performed on a PENTRA® automat. Plasma 25-hydroxyvitamin D was measured by RIA technique and plasma folate and vitamin B₁₂ by competitive immunoassay with direct chemiluminescence detection. Part of the biochemical material was stored at -80°C until its transport to central laboratories for vitamin D, folate and B₁₂ measurements and long-term storage.

Complementary questionnaires. Social and demographic information was collected through face-to-face questionnaires. Previous illnesses such as CVD and diabetes, as well as current drug intake, were described.

Statistical analyses

The theoretical study size was based on recommendations of the EFCOSUM project which advocated performing at least two repeated 24 h recalls in 2000 individuals in order to describe dietary intake at the country level⁽²⁶⁾. Gender-stratified analyses were previously scheduled due to known differences in diet between men and women.

Descriptive analysis of dietary intake comprised subjects with at least two 24 h recalls and basic social and demographic data. Initial weighting was based on the number of eligible individuals multiplied by the inverse of the probability of dwelling selection in the stratum. Calibration was made, separately for each gender, according to the national census data on age, education diploma and whether the household included or did not include at least one child. The final weighting also included the period of data collection. After exclusion of diet-underreporting subjects (see later), a second set of weighting was estimated in subjects who provided a plausible description of their diet. A third weighting set was calculated

for the description of nutritional clinical status in subjects with weight and height available, and a fourth weighting set in subjects with at least the total cholesterol concentration available in order to describe biochemical status. Month of biochemical sample was also taken into account. Means and frequencies with their standard errors were estimated using the STATA® V.10 (Stata Corporation, College Station, TX, USA) declaration for the complex sampling scheme ('svyset' function) and 'svy:' option.

The social and demographic description was made using raw and weighted data in order to evaluate effects of the weighting procedure. Social characteristics were also compared between included and not-included-in-analyses subjects by *t* test and χ^2 tests when appropriate. Factors associated with inclusion at $\alpha < 20\%$ were included in initial multivariate logistic models. The final model included confounding factors and characteristics associated at *P* value <5%. These investigations were performed in subjects included for diet description (*v.* underreporting subjects), anthropometry and blood pressure measurements (*v.* those not included in initial sample) and nutritional markers (*v.* those not included in initial sample and *v.* those included for anthropometry and blood pressure description). Throughout the descriptive analyses, a comparison of male and female data was carried out using test procedures adapted to weighted data.

Dietary intake description. Diet-underreporting subjects were identified by the method proposed by Black⁽²⁷⁾. We used Schofield equations to estimate the BMR according to sex, age and weight (and height when available)⁽²⁸⁾. When weight was missing, BMR was estimated through regression imputation stratified according to age and sex. BMR was compared to energy intake without alcohol (EI) and by taking into account the physical activity level. A physical activity level of 0.88 was used to identify extremely underreporting subjects and a physical activity level of 1.55 to identify other underreporting subjects. Study-internal coefficients were used for the intra-individual energy CV (CV_{wEI}), while those proposed by Black were used for the BMR CV (CV_{wB} = 8.5%) and physical activity CV (CV_{IP} = 15%)⁽²⁷⁾. Comments collected during 24 h recalls were investigated to identify specific conditions that could objectively explain low EI, such as acute disease or low-energy diet to lose weight. Subjects who provided such information were not considered as underreporting subjects, while the other underreporting subjects were excluded from the diet description.

Diet was described as mean energy intake through macronutrients reported on total EI and in grams of food and nutrient intake. Percentages of subjects who attained current national recommendations were estimated. Recommended complex carbohydrate energy (27.5%EI) was based on total carbohydrate intake, added simple sugars and intrinsic sugars. The recommendation for simple sugars from sweetened foods (12.5%) was 10% of added sugars plus 2.5% of intrinsic simple sugars (essentially from milk and fruits). Foods were classified according to the information provided in the PNNS guides⁽¹⁰⁾. Some foods were excluded from a group when the PNNS guides clearly indicated their absence from this group. One dish could pertain to several groups or subgroups if it was composed of several components. Daily mean of each food group intake quantity was computerized in grams. The frequency of consumption was the amount

consumed divided by a portion size which was specific to the type of foods. These frequencies were then compared to PNNS recommendations. When a PNNS recommendation was not clearly quantified⁽¹⁰⁾, international and national recommendations were used.

Physical activity. Data obtained by the IPAQ were computed for a metabolic equivalent task (MET-min/week) following recommendations of the IPAQ group⁽²³⁾. The principle is to multiply the metabolic equivalent task value of each level of physical activity intensity (walking, moderate or vigorous) by the time spent per week. Recommended IPAQ categories of physical activity (low, moderate and high) were used in our analyses⁽²³⁾. Sedentary lifestyle was considered as spending more than 3 h/d in front of a screen (television, computer or videogames).

Nutritional markers. Pregnant women were excluded from the anthropometric description. BMI was calculated as weight in kg/height² in m². The WHO classification was used to define thinness, overweight and obesity, including three grades of obesity⁽²⁴⁾. The European reference for waist circumference was used to define central obesity in men (≥ 94 cm) and women (≥ 80 cm)⁽²⁹⁾. The waist:hip ratio was also calculated; abdominal fat accumulation was estimated through a waist:hip ratio >1.0 in men and >0.85 in women⁽³⁰⁾. The mean of the last two measurements of blood pressure was used in analyses. Hypertension was defined by systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg (French recommendations as in other countries)⁽³¹⁾ and/or intake of drugs affecting blood pressure according to the Anatomical Therapeutic Chemical classification (C02, C03, C07, C08 and C09) (available at <http://www.whocc.no/atcddd/>). Subjects with fasting duration <12 h were excluded from analysis based on lipid profile and glucose. Percentages of adults with lipid components or glucose higher than recommended concentrations in France^(32,33) were described, as well as those taking drugs affecting such components (lipid-lowering drugs (C10) or antidiabetic drugs (A10)). International references were used for anaemia (Hb <130 g/l in men and <120 g/l in women) and for depleted iron stores (ferritin <15.0 μ g/l)⁽³⁴⁾. Finally, references usually cited in the literature were used for low vitamin D status⁽³⁵⁾ and risk of folate deficiency was assessed by plasma folate⁽³⁶⁾.

Results

Participant characteristics

When accounting for withdrawals after initial agreement to participate, the participation rate according to inclusion criteria in analyses for the dietary component of the study was estimated at 59.7% (3115/(4483 eligible households + 734 households in which the contacted person refused to give information on household composition)) (Fig. 1). Missing social and demographic data accounted for 0.8% of subjects having done at least two 24 h recalls. Among subjects who participated in the dietary component of the study, 77.5% were measured for weight and height, and blood samples were taken for 67.5% (Fig. 1).

Mean age, birthplace, marital status, presence of a child at home and smoking habits were unmodified by the weighting




Fig. 1. Inclusion diagram.

procedure (Table 1). Weighted proportions of young adults were higher than raw proportions. Low education level was poorly represented in the sample; primary school level rose about 20% after the weighting procedure. While proportions of service/manual workers and management/intermediate occupations were close in the initial sample, this was no longer the case for weighted proportions (Table 1).

Nutrient and dietary intake

Diet was described in 2734 subjects after exclusion of underreporting subjects (12.2%). The risk of being an underreporting subject increased with increasing age (OR = 1.01, 95% CI 1.00, 1.02, for 1 year), and with place of birth outside France (Europe OR = 1.96, 95% CI 1.14, 3.36; Africa OR = 1.15, 95% CI 0.72, 1.86; other place of birth OR = 3.3, 95% CI 1.56, 6.97, compared to France). Moreover, compared to those who had a primary school level, subjects with a university level were at lower risk of being underreporting subjects (OR = 0.39, 95% CI 0.27, 0.56). Other social and demographic characteristics were comparable between the two groups.

Nutrient intake was higher in men than in women for EI with or without alcohol, complex carbohydrates (%EI), simple sugars (g/d and %EI), total dietary fibre (g/d), calcium

Table 1. Social and demographic characteristics
(Mean values or percentages)

	Raw data		Weighted data*	
	Men	Women	Men	Women
Age				
Mean (years)	46.6	47.0	44.2	45.1
Classes (%)				
18–29	14.9	11.6	22.6	21.5
30–54	51.4	56.2	49.9	49.4
55–74	33.7	32.2	27.5	29.1
Birthplace (%)				
France including overseas territories	90.4	90.5	91.4	91.4
Europe	2.2	3.2	2.9	2.9
Africa	6.4	4.9	5.1	4.2
Others	1.0	1.3	0.5	1.5
Marital status (%)				
Single	19.7	12.4	23.8	17.0
Living with a partner	71.9	70.4	71.6	71.4
Widowed, divorced	8.3	17.2	4.6	11.6
At least one child at home (%)				
Yes	35.9	43.1	35.9	35.5
No	64.1	56.9	64.1	64.5
School level (%)				
Primary school	9.4	13.4	18.4	21.9
Secondary school	35.7	33.8	44.8	40.3
High school	19.8	20.2	17.6	18.5
University	35.1	32.5	19.1	19.3
Occupational category (%)				
Self-employed, farmers	5.9	2.6	5.2	2.4
Management, intermediate professions	30.6	25.0	20.6	16.6
Service worker, manual labour	31.3	34.8	41.2	40.7
Retired	24.4	20.4	21.6	18.1
Home makers, disabled people, others	7.8	17.2	11.5	22.3
Smoking habits (%)				
Never	33.7	53.4	32.2	55.4
Former	35.8	22.3	33.4	18.6
Current	30.5	24.3	34.3	26.0

* Weighting accounted for multilevel sampling design and for social and demographic characteristics compared to the national census (age, school diploma, household including at least one child or not).

(mg/d) ($P < 0.001$ for all tests) and saturated fats (%total lipids; $P = 0.02$) (Table 2). Men also showed a nutrient intake in accordance with recommendations for complex carbohydrates, fibres and calcium that was higher than for women ($P < 0.001$ for all tests) (Table 2). In both sexes, most adults had simple sugar (< 12.5 %EI: 74.5 (SE 1.2) %) and calcium ($\geq 2/3$ French RDA: 76.4 (SE 1.1) %) intake in accordance with recommendations, but compliance with nutrient recommendations was lower for total carbohydrates (> 50 %EI: 26.4 (SE 1.2) %), total fibre (> 25 g/d: 13.7 (SE 0.9) %) and SFA (< 35 % total lipids: 18.5 (SE 1.0) %).

The male diet was characterized by intake of 'bread, cereals and legumes', home-added fats and non-alcoholic beverages mostly in accordance with PNNS recommendations (Table 3). Consumption of fruits and vegetables, seafood and vegetable home-added fat (relatively to animal added fat) was too low compared with PNNS recommendations, while salt intake was too high on average (Table 3). Less than one-third of men had dairy product intake in accordance with the PNNS guidelines (Table 3) and 25.8 (SE 1.8) % had a higher intake than the PNNS guidelines. With an intake of 383 g/d on average, consumption of fruits and vegetables was 400 g/d or higher in 44 % of men. Alcohol consumption

was higher than the recommendation for 20.0 (SE 1.7) % of men. In addition, 25.6 (SE 1.9) % had salt intake higher than 12 g/d. In women, 'meat, seafood and egg' intake was adequate for most of them, as was home-added fat, sweetened and alcoholic beverages, and salt (Table 3). Dairy product intake was lower than the PNNS guidelines in 54.6 (SE 1.6) % of them, while about one-quarter had satisfactory consumption. Intake of 'bread, cereals and legumes' was also much too low (Table 3), with 59.6 (SE 1.5) % of women consuming less than the PNNS guidelines. Finally, more than half of women had fruit and vegetable intake lower than 400 g/d.

Physical activity and sedentary lifestyle

Behaviours were described in 2971 subjects due to the fact that there were 4.6 % 'I don't know'-type answers along with missing data on physical activity. Social and demographic characteristics were comparable between subjects included and not included in physical activity analysis. According to the IPAQ classification, 36.8 (SE 1.2) % of adults had a low level of physical activity, 36.7 (SE 1.1) % an intermediate level and 26.5 (SE 1.1) % had a high level of physical activity. Low physical activity was comparable

Table 2. Daily mean energy and nutrient intake and percentage of adherence to guidelines*

(Mean values or percentages with their standard errors)

	Men		Women	
	Mean or %	SE	Mean or %	SE
Mean energy intake				
Without alcohol (MJ/d)	9.47	0.1	6.96	0.1
Without alcohol (kcal/d)	2264	26	1664	14
Including alcohol (MJ/d)	9.97	0.1	7.10	0.1
Including alcohol (kcal/d)	2384	28	1698	14
% EI from carbohydrate				
Total carbohydrate				
Mean (%EI)	45.8	0.3	45.5	0.2
% > 50%EI	28.1	2.0	24.7	1.4
Complex carbohydrate				
Mean (%EI)	26.8	0.3	24.3	0.2
% ≥ 27.5%EI	44.1	2.1	27.8	1.4
Simple sugar				
Mean (g/d)	109	2	87	1
Mean (%EI)	19.0	0.3	21.1	0.2
% < 12.5%EI	74.1	1.9	74.9	1.5
Total dietary fibre†				
Mean (g/d)	19.1	0.3	15.9	0.2
% > 25 g/d	19.3	1.6	8.3	0.8
% EI from fat				
Total fat				
Mean (%EI)	37.1	0.2	37.1	0.2
% < 35%EI	35.7	2.0	37.7	1.6
SFA				
Mean (% total lipids)	40.6	0.2	39.9	0.2
% < 35% total lipids	17.0	1.6	20.0	1.3
Calcium				
Mean (mg/d)	974	14	837	11
% < 2/3 RDA‡	15.4	1.6	31.6	1.5

EI, energy intake without alcohol.

* For details of subjects and procedures, see Methods and Table 1.

† American Association of Analytical Chemists method.

‡ French RDA: 900 mg/d in adults > 18 years, excepting 1200 mg/d in women > 55 years and men > 65 years.

in men and women, but the proportion of men with a high level of physical activity was higher than that of women ($P=0.02$) (Table 4).

Time spent in front of a screen was about 3 h 20 min/d, which was overall much higher in men than in women ($P<0.001$) (Table 4). During working days, such sedentary activity (mean duration: 161 (SE 3.3) min/d) was, as expected, less long than during leisure days (mean duration: 222 (SE 3.9) min/d). During working days, television viewing time was comparable between men and women, but not during leisure days ($P=0.03$) (Table 4). Finally, some two-thirds of adults spent more than 2 h/d watching television, whatever the day.

Clinical and biochemical markers of nutritional status

Anthropometry and blood pressure were described in 2413 adults. These subjects were not comparable to those not included ($n=702$) for age (OR = 1.02, 95% CI 1.01, 1.03, for 1 year), living status (living with a partner OR = 1.32, 95% CI 1.04, 1.68, compared to being single) or school level (secondary school OR = 1.37, 95% CI 1.02, 1.84; high school OR = 1.40, 95% CI 1.01, 1.95; university OR = 1.85, 95% CI 1.36, 2.52, compared to primary school). Biochemical

markers were described in 2102 subjects who were older (OR = 1.02, 95% CI 1.01, 1.02, for 1 year) and had a higher education level (secondary OR = 1.48, 95% CI 1.14, 1.92; high school OR = 1.39, 95% CI 1.04, 1.86; university OR = 2.11, 95% CI 1.60, 2.78, compared to primary school level) than those not included ($n=1013$). In the end, subjects included in the biochemical description were different from those included in anthropometry and blood pressure descriptions in terms of sex (women OR = 0.72, 95% CI 0.55, 0.95, compared to men) or school level (university OR = 2.06, 95% CI 1.38, 3.08, compared to primary school).

About half of adults were found to be at high cardiovascular risk based on hypercholesterolaemia and intake of lipid-lowering treatments (61.4 (SE 1.6) %), overweight (49.3 (SE 1.4) %) and dyslipidaemia (44.1 (SE 1.6) %). Frequencies of high waist:hip ratio (21.1 (SE 1.2) %) and hypertension (30.9 (SE 1.3) %) were lower. Despite comparable obesity frequency, women were more frequently at cardiovascular risk than men, based on waist circumference ($P=0.002$), waist:hip ratio ($P<0.001$) and elevated cholesterol (not taking into account drug intake; $P=0.04$) (Table 5). In contrast, men were more frequently affected by high blood pressure ($P<0.001$), hypertension ($P=0.02$), dyslipidaemia ($P<0.001$), hyperglycaemia ($P=0.015$) and diabetes ($P=0.015$) (Table 5).

Iron deficiency was more frequent in women than in men ($P=0.03$) (Table 6). In women of childbearing age (18–49 years of age), frequency of iron deficiency was 3.0 (SE 0.7) %, frequency of anaemia was 5.7 (SE 1.0) %, and 13.5 (SE 1.8) % of them had depleted iron stores based on concentrations of plasma ferritin. Risk of folate deficiency was observed in 6.2 (SE 0.9) % of adults. Mean plasma folate was higher in women than in men ($P<0.001$), but the frequency risk of deficiency was comparable (Table 6). A total of 7.7 (SE 1.7) % of women of childbearing age presented such a risk. Low vitamin D status was revealed in 4.5% (SE 0.6) of adults, with higher risk in women than in men ($P=0.04$) (Table 6). Finally, low status of plasma vitamin B₁₂ was observed in 3.3 (SE 0.6) % of adults.

Discussion

The ENNS survey provides a recent overview on dietary intake, physical activity and nutritional status in 18–74-year-old adults in France. The diet appears to be heterogeneous, with satisfactory intakes of total energy, simple sugar, calcium and alcohol, but does not have a well-balanced intake of macronutrient, SFA or total fibre according to current recommendations. While fruit and vegetable consumption and fish intake were lower than current PNNS guidelines, food intake was generally in compliance with recommendations for other food groups. However intake of salt and dairy products was too high in men. In contrast, 'bread, cereals and legumes' and dairy product consumption was below the guidelines in women. According to anthropometry, blood pressure and lipid profiles, cardiovascular risk was elevated in 18–74-year-old adults. In other respects, vitamin status was at risk of deficiency in less than 10% of adults. Finally, over two-third of adults had satisfactory physical activity, but most spent a long time in front of a screen.

The present survey, the first of its kind to be conducted in France, is of interest in surveillance and monitoring of

Table 3. Daily mean diet intake and percentages of adherence to guidelines*
(Mean values or percentages with their standard errors)

	Men		Women	
	Mean or %	SE	Mean or %	SE
Fruits and vegetables†				
Mean (g/d)	383	10	389	6
% ≥ 400 g/d	43.7	2.1	43.9	1.6
Dairy products‡				
Mean (g/d)	214	6.9	212	4.8
% 2.5–3.5 servings/d	30.2	1.9	28.1	1.4
Meat, seafood and eggs (excluding processed meat)				
Mean (g/d)	166	3	126	2
% 1–2 servings/d	50.3	2.1	54.6	1.6
Seafood ≥ 2 servings/week	27.7	1.8	32.1	1.5
Bread, cereals, legumes, potatoes				
Mean (g/d)	337	5	230	3
% 3–6 servings/d	60.1	2.1	38.2	1.5
Home-added fat (butter, oils, margarine)				
Mean (g/d)	33.0	0.8	29.1	0.6
% < 16% EI	87.0	1.5	81.9	1.2
% > 50% vegetable/total added fat ratio	41.2	2.0	47.6	1.6
Non-alcoholic or added sweetened beverage				
Mean (l/d)	1.60	0.03	1.49	0.02
% < 250 ml/d of sweetened beverage	84.8	1.7	92.3	1.1
Alcoholic beverages				
Non-consumers (%)	13.2	1.6	27.3	1.5
% < 2 servings/d in women and 3 servings/d in men§	66.9	2.1	66.5	1.5
Total salt				
Mean (g/d)	9.9	0.1	7.0	0.1
% < 6 g/d	8.7	1.2	35.3	1.6
% < 8 g/d	29.8	2.0	69.8	1.4

EI, energy intake without alcohol.

* For details of subjects and procedures, see Methods and Table 1. For each food group, components of composite dishes were proportionally accounted for.

† Including 100% juices, processed fruit and vegetables; excluding legumes, beans, starchy plants and potatoes.

‡ Excluding desserts with high contents of sugar and fat. Range in 55–74-year-old adults: 2.5–4.5 servings/d.

§ One serving is equivalent to a glass of wine, i.e. 10 g alcohol; only includes consumers.

|| Including salt added when cooking and during meals; estimated using 24 h recalls.

public health programmes. In the present study, the participation rate in diet data collection attained 60%, considered statistically acceptable, but the overall participation rate fell to 40% for biochemical sampling. Improvement in the participation rate is a critical point in such studies so as to minimize the selection bias as much as possible. It is unknown as to whether initial contact, organized home visits, survey topics, individual and global result return, and staff quality had an effect upon the participation rate⁽³⁷⁾. No incentive was offered to subjects but participation in the health examination was expanded by home visits from a nurse. Despite a high acceptability rate and quality of the data, the home visit strategy is costly⁽³⁸⁾. Comparisons of crude and weighting data as well as subject characteristics according to their entire or partial participation showed recruitment biases. In particular, low-income subjects were less often included than would be desirable. These biases were compensated for by weighting statistics.

Nutrient and dietary intake

Comparisons to Western nutritional surveys are limited due to variations in sampling schemes and dietary collection methods. Data treatment also has an impact on observations (food group classification, nutritional value table). However, the diet

description shows patterns comparable to previous descriptive Western studies, including insufficient carbohydrate and elevated fat intake, especially that of saturated fats⁽¹³⁾. Recent US trends show a decrease in fat energy intake along with an increase in total energy and carbohydrate intake⁽¹¹⁾. Repeated European surveys do not clearly indicate whether such changes are ongoing or not^(16,39). Trends are difficult to interpret in France due to the absence of published studies using comparable methods⁽¹⁷⁾. Salt intake was generally too high, especially in men, although it seemed to progressively attain current French recommendations. As previously shown⁽¹⁴⁾, this might be related to a decrease in energy intake.

These patterns are consistent with food group consumption described as complying with PNNS recommendations. The insufficient calcium intake in women is partly related to insufficient consumption of dairy products and milk. In France, lipid intake seems to depend more on consumption of foods that contain fat such as processed meat, cheese, snacking foods etc. rather than on home-added fats such as butter, margarine, oils etc. Indeed, while the lipid intake remained too high, home-added fat consumption was satisfactory. Regarding food groups globally, the French nutritional situation would appear to be intermediate between a protective Mediterranean diet in southern Europe and a northern European diet⁽⁴⁰⁾. Variations had also been observed within the French

Table 4. Physical activity and sedentary lifestyle indicators*
(Mean values or percentages with their standard errors)

	Men		Women	
	Mean or %	SE	Mean or %	SE
Physical activity				
Mean MET-min/week	2679	102	2130	63
IPAQ categories (%)				
Low	36.1	2.0	37.5	1.4
Intermediate	34.4	1.9	38.9	1.5
High	29.5	1.8	23.6	1.3
Mean time in front of a screen (television, computer, videogames)				
Daily mean†				
Mean time (min/d)	216	5	186	3
% ≥ 180 min/d	58.8	2.0	47.9	1.5
Working days				
Mean time (min/d)	173	5	147	4
% ≥ 180 min/d	47.8	2.5	32.9	2.0
Leisure days				
Mean time (min/d)	243	7	201	4
% ≥ 180 min/d	64.7	2.0	55.5	1.5
Mean time in front of a television				
Daily mean†				
Mean time (min/d)	161	4	155	3
% ≥ 120 min/d	67.1	1.9	66.8	1.4
Working days				
Mean time (min/d)	120	4	117	3
% ≥ 120 min/d	59.2	2.5	54.8	2.1
Leisure days				
Mean time (min/d)	181	5	168	3
% ≥ 120 min/d	73.7	1.9	74.0	1.3

IPAQ, International Physical Activity Questionnaire; MET, metabolic equivalent task.

* For details of subjects and procedures, see Methods and Table 1.

† Weighted mean of working days and leisure days.

territory up until recently⁽¹⁹⁾. Regional variations were taken into account in our survey, by recruiting throughout continental France.

The dietary overview provided by the ENNS survey must be interpreted with caution, since some limitations were present. Three 24 h recalls were carried out in a 15 d period: at the individual level, this is not sufficient to describe the usual intake. The description of some nutrient intakes (carbohydrates and calcium, for instance) might require more than three 24 h recalls⁽⁴¹⁾. However, the present data are reliable at the group level, since recruitment and diet information collection were carried out throughout the entire territory during a 1-year period⁽²⁶⁾. For calcium intake, supplements are not taken into account, since PNNS guidelines explicitly discourage their consumption, with a few specific exceptions⁽¹⁰⁾.

Physical activity and sedentary lifestyle

Regarding physical activity, one European survey, using the short-form IPAQ, showed that the prevalence of satisfactory activity (e.g. the high category of IPAQ) in France (24.1%) was among the lowest frequencies in Europe, while frequencies in Germany (40.2%) and the Netherlands (44.2%) were among the highest⁽⁴²⁾. The difference between men and women might be explained by leisure-time physical activity, especially sports, more commonly practised by men than women in France⁽⁴³⁾. Despite conclusive validation⁽²³⁾, IPAQ

use is considered to lead to overestimation of moderate and vigorous physical activity. The quality of the interview is crucial and must be carefully prepared. Indeed, it is a useful tool for surveillance, since physical activity is examined under all conditions (work, transportation, leisure time, etc.)

Clinical and biochemical markers of nutritional status

The French dietary pattern, along with insufficient physical activity, helps to explain the high prevalence of cardiovascular risk observed in the ENNS survey. In France, body mass changes have been mainly monitored during the last decade by surveys using self-measured weight and height⁽¹⁸⁾. Between 1997 and 2006, a continuous increase in overweight and obesity has been observed in both male and female adults (obesity prevalence equal to 12.5 and 13.6%, respectively, in 2006). As expected⁽⁴⁴⁾, we found a higher cross-sectional prevalence of obesity based on measured anthropometry. Based on declared anthropometry, the French frequency of overweight and obesity was among the lowest in Europe up until the beginning of the 2000s⁽⁴⁵⁾, but comparisons should now be made using measured anthropometry. For example, when compared to the 14.2% obesity based on measured data in 2003–5 in 18–64-year-old Portuguese adults⁽⁴⁶⁾, the French prevalence was somewhat higher.

Using comparable methods of measurement but different age categories (18–74 years *v.* ≥ 18 years), the prevalence of hypertension in 2006–7 in France was equal to the frequency observed in 2003–4 in the USA (29.6%)⁽¹²⁾. Compared to European data, mean systolic blood pressure in France in 2001 was comparable to the mean reported in 19–64-year-old adults in the UK (130 mm Hg in men and 122 mm Hg in women), whereas this was not the case for diastolic blood pressure (73 and 68 mm Hg, respectively)⁽⁴⁷⁾. Methods of measurements must be taken into account when interpreting variations across surveys.

In France, adults appeared to be exposed to factors leading to metabolic syndrome, diabetes and their associated cardiovascular manifestations, but differed from other Western countries. For example, the mean total cholesterol and frequency of elevated total cholesterol were higher than reported values from the 1999–2004 National Health and Nutrition Examination Survey but with a frequency of high LDL-cholesterol comparable in women and lower in men⁽⁴⁸⁾. The same tendency was observed when comparison was made with similar surveys in the UK⁽⁴⁷⁾ and in Finland⁽⁴⁹⁾. This illustrates the widely described 'French paradox' of elevated total cholesterol and dyslipidaemia but lower LDL-cholesterol and higher HDL-cholesterol concentrations. Despite an insufficient explanation *per se*, such variations could contribute to understanding variations in cardiovascular mortality across Europe⁽⁷⁾.

For biomarkers of vitamin and mineral status, comparable surveys aimed at describing overall nutritional status are lacking, except for the National Diet and Nutrition Survey conducted in 2001–2 in the UK⁽⁴⁷⁾. One of our limitations compared to the UK survey was that Hb and ferritin were measured in field laboratories, while UK analyses were centralized. However, folate, vitamin B₁₂ and vitamin D measurements were centralized in the ENNS survey, the assay methods being comparable with those used in the UK survey. Anaemia and serum ferritin depletion were more frequent in

Table 5. Anthropometry, blood pressure, cholesterol and glucose status*
(Mean values or percentages with their standard errors)

	Men		Women	
	Mean or %	SE	Mean or %	SE
Anthropometric measurement†				
Mean weight (kg)	79.2	0.6	65.6	0.5
Mean height (m)	174.4	0.3	161.7	0.2
Mean hip circumference (cm)	99.6	0.5	100.7	0.4
Waist circumference (cm)				
Mean	92.3	0.6	81.9	0.4
% M: ≥ 94 cm; W: ≥ 80 cm	42.3	2.3	51.6	1.8
Waist:hip ratio†				
Mean	0.93	0.005	0.81	0.003
% M: > 1.0 ; W: > 0.85	14.6	1.7	27.6	1.5
BMI†				
Mean (kg/m ²)	26.1	0.2	25.1	0.2
% Thinness (BMI < 18.5)	0.7	0.3	5.0	0.8
% Overweight (BMI ≥ 25.0)	57.2	2.3	41.4	1.7
% Obesity (BMI ≥ 30.0)	16.1	1.6	17.6	1.3
Grade I (30.0 \leq BMI \leq 34.9)	12.9	1.4	12.1	1.1
Grade II (35.0 \leq BMI \leq 39.9)	2.5	0.8	4.2	0.8
Grade III (40.0 \leq BMI)	0.7	0.3	1.2	0.3
BP				
Mean systolic BP (mm Hg)	129	0.8	119	0.6
Mean diastolic BP (mm Hg)	79	0.6	76	0.4
% Elevated BP‡	28.3	2.1	17.8	1.3
% Hypertension§	34.1	2.2	27.8	1.5
Fasting serum cholesterol				
Total cholesterol				
Mean (mmol/l)	5.31	0.06	5.44	0.04
% ≥ 5.16 mmol/l	52.6	2.5	59.0	1.9
% ≥ 5.16 mmol/l and/or lipid-lowering drugs	60.7	2.5	62.1	1.9
LDL-cholesterol (mmol/l)				
Mean (mmol/l)	3.31	0.05	3.29	0.04
% ≥ 4.10 mmol/l	22.3	2.2	18.5	1.4
% ≥ 4.10 mmol/l and/or lipid-lowering drugs	34.8	2.4	27.9	1.7
% Dyslipidaemia	50.9	2.5	37.6	1.9
Fasting serum glucose				
Mean (mmol/l)	5.42	0.06	5.07	0.04
% ≥ 7.0 mmol/l	5.2	1.1	2.5	0.6
% ≥ 7.0 mmol/l and/or anti-diabetic drugs¶	6.1	1.1	3.2	0.6

BP, blood pressure; M, men; W, women.

* For details of subjects and procedures, see Methods and Table 1.

† Pregnant women excluded (*n* 25).‡ Systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mm Hg.

§ Elevated BP and/or taking drugs acting on BP.

|| TAG ≥ 1.7 mmol/l and/or LDL-cholesterol ≥ 4.1 mmol/l and/or HDL-cholesterol ≤ 1.0 mmol/l and/or lipid-lowering drug intake.

¶ Non-fasting in subjects taking anti-diabetic drugs.

France, especially in women, than in the UK. Folate status was also less satisfactory in France than in the UK, possibly due to supplements and enriched foods, which may be less frequently consumed in France. The vitamin B₁₂ status was comparable, however, in the two surveys. In contrast, and as expected, the vitamin D status was much more satisfactory in France than in the UK, where about 14 % of adults⁽⁴⁷⁾ were at risk of deficiency compared to 4.5 % in France. Variations in diet between the two countries should be carefully examined in order to understand these results.

The ENNS survey provides a complete description of nutritional characteristics in adults using reference methods recommended for diet, physical activity and nutritional status investigation. The present study elucidates a contrasting but globally unsatisfactory nutritional situation in France. This data set constitutes an essential tool for monitoring public health pro-

grammes such as the PNNS. Beyond national monitoring, such data are useful for international comparisons in order to better understand morbidity and mortality variations across countries.

Acknowledgements

ENNS was supported by the French Institut de veille sanitaire (InVS), the Université de Paris 13, and the Caisse nationale d'assurance maladie des travailleurs salariés (CnamTS). We thank the dietitians, nurses and physicians, the staff of Health Examination Centres of CnamTS, their laboratories, the Grenoble University Hospital Center (Département de Biologie Intégrée), U.631 Inserm and the Pasteur Institute which contributed to data collection and biochemical analyses. K. C. designed the study, managed the data acquisition,

Table 6. Nutritional biomarkers*
(Mean values or percentages with their standard errors)

	Men		Women	
	Mean or %	SE	Mean or %	SE
Iron status				
Hb				
Mean (g/l)	153	0.5	137	0.4
% M: <130 g/l; W: <120 g/l	1.9	0.6	4.5	0.7
Serum ferritin				
Mean ($\mu\text{g/l}$)	291	93	81	3
% <15.0 $\mu\text{g/l}$	1.3	0.5	8.7	1.2
% 15.0–30.0 $\mu\text{g/l}$	2.6	1.0	18.0	1.6
Iron-deficiency status (%)				
W: Hb <120 g/l and ferritin <15.0 $\mu\text{g/l}$			1.9	0.4
M: Hb <130 g/l and ferritin <15.0 $\mu\text{g/l}$	0.5	0.3		
Serum 25-hydroxyvitamin D				
Mean (nmol/l)	59.8	1.2	56.4	1.1
% <25 nmol/l	3.3	0.7	5.5	0.9
Plasma folate				
Mean (nmol/l)	14.2	0.3	16.4	0.3
% <7 nmol/l	6.7	1.3	5.8	1.1
Plasma vitamin B ₁₂				
Mean (pmol/l)	289	4	304	6
% <150 pmol/l	2.9	0.8	3.7	0.8

M, men; W, women.

*For details of subjects and procedures, see Methods and Table 1.

performed statistical analysis, interpreted the data and wrote the manuscript. M. V., A. M., B. S. and V. D. contributed to the management of data collection, performed the statistical analysis, interpreted the data and revised the manuscript. C. R., A. O. and E. S. contributed to the data collection, provided administrative and material support, and participated in the statistical analysis. S. H. designed the study, contributed to the interpretation of results and revised the manuscript. All authors read and approved the final version of the manuscript. All authors declare no personal or financial interest in the content of the paper.

References

- World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Joint WHO/FAO Expert Consultation. WHO Technical Report Series*, no. 916. Geneva: WHO.
- Dalziel K & Segal L (2007) Time to give nutrition interventions a higher profile: cost-effectiveness of 10 nutrition interventions. *Health Promot Int* **22**, 271–283.
- van Baal PH, Polder JJ, de Wit GA, *et al.* (2008) Lifetime medical costs of obesity: prevention no cure for increasing health expenditure. *PLoS Med* **5**, e29.
- Lachat C, Van Camp J, de Henauw S, *et al.* (2005) A concise overview of national nutrition action plans in the European Union Member States. *Public Health Nutr* **8**, 266–274.
- Emery C, Dinet J, Lafuma A, *et al.* (2007) Evaluation du coût associé à l'obésité en France (Cost of obesity in France). *Presse Med* **36**, 832–840.
- Belot A, Grosclaude P, Bossard N, *et al.* (2008) Cancer incidence and mortality in France over the period 1980–2005. *Rev Epidemiol Sante Publique* **56**, 159–175.
- Muller-Nordhorn J, Binting S, Roll S, *et al.* (2008) An update on regional variation in cardiovascular mortality within Europe. *Eur Heart J* **29**, 1316–1326.
- Kusnik-Joinville O, Weill A, Salanave B, *et al.* (2008) Prevalence and treatment of diabetes in France: trends between 2000 and 2005. *Diabetes Metab* **34**, 266–272.
- Samson S, Ricordeau P, Pépin S, *et al.* (2007) Hypertension artérielle et facteurs de risque associés: évolutions des traitements entre 2000 et 2006 (Hypertension and associated risk factors: changes in drugs between 2000 and 2006). *Points de Repère (CnamTS)* **10**, 1–8.
- Hercberg S, Chat-Yung S & Chauillac M (2008) The French national nutrition and health program: 2001–2006–2010. *Int J Public Health* **53**, 68–77.
- Briefel RR & Johnson CL (2004) Secular trends in dietary intake in the United States. *Annu Rev Nutr* **24**, 401–431.
- Ong KL, Cheung BM, Man YB, *et al.* (2007) Prevalence, awareness, treatment, and control of hypertension among United States adults 1999–2004. *Hypertension* **49**, 69–75.
- Elmadfa I, Weichselbaum E, König J, *et al.* (2005) European nutrition and health report 2004. *Forum Nutr* **58**, 1–220.
- Reinivuo H, Valsta LM, Laatikainen T, *et al.* (2006) Sodium in the Finnish diet: II trends in dietary sodium intake and comparison between intake and 24-h excretion of sodium. *Eur J Clin Nutr* **60**, 1160–1167.
- Swan G (2004) Findings from the latest National Diet and Nutrition Survey. *Proc Nutr Soc* **63**, 505–512.
- Turrini A, Saba A, Perrone D, *et al.* (2001) Food consumption patterns in Italy: the INN-CA Study 1994–1996. *Eur J Clin Nutr* **55**, 571–588.
- Volatier JL & Verger P (1999) Recent national French food and nutrient intake data. *Br J Nutr* **81**, Suppl. 2, S57–S59.
- Charles MA, Eschwege E & Basdevant A (2008) Monitoring the obesity epidemic in France: the Obepi surveys 1997–2006. *Obesity (Silver Spring)* **16**, 2182–2186.
- Kesse E, Boutron-Ruault MC & Clavel-Chapelon F (2005) Regional dietary habits of French women born between 1925 and 1950. *Eur J Nutr* **44**, 285–292.
- Perrin AE, Dallongeville J, Ducimetiere P, *et al.* (2005) Interactions between traditional regional determinants

- and socio-economic status on dietary patterns in a sample of French men. *Br J Nutr* **93**, 109–114.
21. Le Moulllec N, Deheeger M, Preziosi P, *et al.* (1996) Validation du manuel-photos utilisé pour l'enquête alimentaire de l'étude SU.VI.MAX (Validation of the photography manual of servings used in dietary collection in the SU.VI.MAX study). *Cah Nutr Diet* **31**, 158–164.
 22. Herceberg S, (ed). (2005) *Table de Composition des Aliments – SU.VI.MAX*. Paris: Economica.
 23. Craig CL, Marshall AL, Sjostrom M, *et al.* (2003) International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* **35**, 1381–1395.
 24. World Health Organization (1995) *Physical Status: The Use and Interpretation of Anthropometry. Report of a WHO Expert Committee*. Geneva: WHO.
 25. Friedewald WT, Levy RI & Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* **18**, 499–502.
 26. Brussaard JH, Lowik MR, Steingrimsdottir L, *et al.* (2002) A European food consumption survey method – conclusions and recommendations. *Eur J Clin Nutr* **56**, Suppl. 2, S89–S94.
 27. Black AE (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes Relat Metab Disord* **24**, 1119–1130.
 28. Schofield WN (1985) Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* **39**, Suppl. 1, 5–41.
 29. Balkau B & Charles MA (1999) Comment on the provisional report from the WHO consultation. European Group for the Study of Insulin Resistance (EGIR). *Diabet Med* **16**, 442–443.
 30. World Health Organization (1998) *Preventing and Managing the Global Epidemic. Report of a WHO Consultation on Obesity*. Geneva: WHO.
 31. Pickering TG, Hall JE, Appel LJ, *et al.* (2005) Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension* **45**, 142–161.
 32. Agence Française de Sécurité Sanitaire des Produits de Santé (Afssaps) (2005) *Prise en charge thérapeutique du patient dyslipidémique (Care of Dyslipidemic Patients)*. Paris: Afssaps.
 33. Haute Autorité de Santé (2005) *Prise en charge des patients adultes atteints d'hypertension artérielle essentielle – Actualisation 2005 – Recommandations (Care of Essential Hypertension – 2005 Update and Recommendations)*. Paris: HAS.
 34. UNICEF, UNU & WHO (2001) *Iron Deficiency Anaemia: Assessment, Prevention, and Control. A Guide for Programme Managers no. WHO/NHD/01.3*. Geneva: WHO.
 35. Lips P (2004) Which circulating level of 25-hydroxyvitamin D is appropriate? *J Steroid Biochem Mol Biol* **89/90**, 611–614.
 36. Jacques PF, Selhub J, Bostom AG, *et al.* (1999) The effect of folic acid fortification on plasma folate and total homocysteine concentrations. *N Engl J Med* **340**, 1449–1454.
 37. Mapstone J, Elbourne D & Roberts IG (2007) Strategies to improve recruitment to research studies. *The Cochrane Database of Systematic Reviews 2007*, issue 2, MR000013. <http://www.mrw.interscience.wiley.com/cochrane/clsysrev/articles/MR000013/frame.html>.
 38. Siemiatycki J (1979) A comparison of mail, telephone, and home interview strategies for household health surveys. *Am J Public Health* **69**, 238–245.
 39. Serra-Majem L, Ribas-Barba L, Salvador G, *et al.* (2007) Trends in energy and nutrient intake and risk of inadequate intakes in Catalonia, Spain (1992–2003). *Public Health Nutr* **10**, 1354–1367.
 40. Slimani N, Fahey M, Welch AA, *et al.* (2002) Diversity of dietary patterns observed in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* **5**, 1311–1328.
 41. Mennen LI, Bertrais S, Galan P, *et al.* (2002) The use of computerized 24 h dietary recalls in the French SU.VI.MAX study: number of recalls required. *Eur J Clin Nutr* **56**, 659–665.
 42. Sjostrom M, Oja P, Hagstromer M, *et al.* (2006) Health-enhancing physical activity across European Union countries: the Eurobarometer study. *J Public Health (Springer)* **14**, 291–300.
 43. Oppert JM, Thomas F, Charles MA, *et al.* (2006) Leisure-time and occupational physical activity in relation to cardiovascular risk factors and eating habits in French adults. *Public Health Nutr* **9**, 746–754.
 44. Niedhammer I, Bugel I, Bonenfant S, *et al.* (2000) Validity of self-reported weight and height in the French GAZEL cohort. *Int J Obes* **24**, 1111–1118.
 45. Seidell JC (2002) Prevalence and time trends of obesity in Europe. *J Endocrinol Invest* **25**, 816–822.
 46. doCarmo I, DosSantos O, Camolas J, *et al.* (2008) Overweight and obesity in Portugal: national prevalence in 2003–2005. *Obes Rev* **9**, 11–19.
 47. Ruston D, Hoare J, Henderson L, *et al.* (2004) *The National Diet & Nutrition Survey: Adults Aged 19 to 64 Years – Nutritional Status (Anthropometry and Blood Analytes), Blood Pressure and Physical Activity*, vol. 4. London: Food Standards Agency.
 48. Hyre AD, Muntner P, Menke A, *et al.* (2007) Trends in ATP-III-defined high blood cholesterol prevalence, awareness, treatment and control among U.S. adults. *Ann Epidemiol* **17**, 548–555.
 49. Ilanne-Parikka P, Eriksson JG, Lindstrom J, *et al.* (2004) Prevalence of the metabolic syndrome and its components: findings from a Finnish general population sample and the Diabetes Prevention Study cohort. *Diabetes Care* **27**, 2135–2140.