

## Iodine Content of Prenatal Multivitamins in the United States

**TO THE EDITOR:** Normal thyroid function in fetuses and breast-fed infants, which is dependent on sufficient maternal dietary intake of iodine, is crucial for normal neurocognitive development.<sup>1</sup> Children of mothers with severe iodine deficiency may have cretinism. Iodine deficiency affects more than 2.2 billion persons (38% of the world's population) and is the leading cause of preventable mental retardation worldwide.<sup>2</sup> Even mild iodine deficiency may have adverse effects on the cognitive function of children.<sup>1</sup>

National surveys including the first and third National Health and Nutrition Examination Surveys (NHANES I and III) and NHANES 2001–2002 have reported a decrease of approximately 50% in adult urinary iodine values since the 1970s in the United States. In the subgroup of women of child-bearing age, the median urinary iodine value decreased from 294 to 128  $\mu\text{g}$  per liter. The most recent NHANES survey (2003–2004) reported that 37.2% of this subgroup of women had urinary iodine values below 100  $\mu\text{g}$  per liter, which suggests mild iodine deficiency.<sup>3</sup>

The Institute of Medicine recommends a daily iodine intake of 220  $\mu\text{g}$  during pregnancy and 290  $\mu\text{g}$  during lactation; the World Health Organization recommends 250  $\mu\text{g}$  of iodine daily for pregnant and lactating women. The American Thyroid Association has recommended that women receive prenatal vitamins containing 150  $\mu\text{g}$  of iodine daily during pregnancy and lactation.<sup>4</sup> However, the iodine content of prenatal multivitamins is not mandated in the United States.

Using the Internet, we identified 127 nonprescription and 96 prescription prenatal multivitamins currently marketed in the United States. Of these multivitamins, 114 (87 nonprescription and 27 prescription) contained iodine; according to the product labeling, 101 (89%) contained 150  $\mu\text{g}$  or more of iodine per daily dose. The iodine was in the form of kelp (in 42 multivitamins), potassium iodide (in 67), or another ingredient (in 5).

We measured the iodine content in 60 randomly selected iodine-containing prenatal multivitamins (1 to 2 tablets per bottle, each from a single lot) and compared the results with the values on their labels (Table 1). The mean ( $\pm\text{SE}$ ) level of measured iodine per daily dose in 35 potassium iodide-containing vitamins was  $119.0\pm 13.6$   $\mu\text{g}$ .

**Table 1. Iodine Content in Prenatal Multivitamins.**

Multivitamins Derived from Potassium Iodide (N=35)		Multivitamins Derived from Kelp (N=25)	
Iodine Value on Label	Measured Iodine	Iodine Value on Label	Measured Iodine
$\mu\text{g}/\text{daily dose}$		$\mu\text{g}/\text{daily dose}$	
Nonprescription		Nonprescription	
25	11	75	55
50	49	150	33
75	15	150	36
150	17	150	40
150	31	150	55
150	36	150	67
150	91	150	70
150	95	150	70
150	100	150	75
150	107	150	115
150	114	150	115
150	152	150	132
150	228	150	144
300	213	150	155
300	236	150	180
300	395	150	214
Prescription		150	240
150	26	150	270
150	46	180	198
150	49	200	115
150	56	225	110
150	87	225	227
150	91	225	229
150	103	226	610
150	106	300	46
150	116		
150	117		
150	117		
150	129		
150	144		
150	144		
150	162		
150	167		
150	220		
175	186		
200	205		

Potassium iodide contains 76% iodide. Thus, the measured iodine content was approximately equivalent to 76% of the total potassium iodide content. The measured iodine in 25 brands containing kelp was 33 to 610  $\mu\text{g}$  per daily dose. Thirteen brands contained levels of iodine that were discordant by 50% or more with the values on their labels, including 10 brands with iodine values that were lower by 50% or more. Variations in the iodine content of kelp have been reported previously.<sup>5</sup>

Manufacturers of prenatal multivitamins in the United States should be encouraged to use only potassium iodide, to maintain consistency in labeling, and to ensure that these vitamins contain 150  $\mu\text{g}$  of supplemental daily iodine by including at least 197  $\mu\text{g}$  of potassium iodide per daily dose, as recommended by the American Thyroid Association.

Angela M. Leung, M.D.

Elizabeth N. Pearce, M.D.

Lewis E. Braverman, M.D.

Boston University Medical Center  
Boston, MA 02118  
elizabeth.pearce@bmc.org

1. de Escobar DM, Obregón MJ, del Rey FE. Maternal thyroid hormones early in pregnancy and fetal brain development. *Best Pract Res Clin Endocrinol Metab* 2004;18:225-48.
2. Iodine deficiency. Ottawa: International Council for the Control of Iodine Deficiency Disorders. (Accessed February 5, 2009, at <http://www.iccid.org/pages/iodine-deficiency.php>.)
3. Caldwell KL, Miller GA, Wang RY, Jain RB, Jones RL. Iodine status of the U.S. population, National Health and Nutrition Examination Survey 2003–2004. *Thyroid* 2008;18:1207-14.
4. Becker DV, Braverman LE, Delange F, et al. Iodine supplementation for pregnancy and lactation — United States and Canada: recommendations of the American Thyroid Association. *Thyroid* 2006;16:949-51.
5. Teas J, Pino S, Critchley A, Braverman LE. Variability of iodine content in common commercially available edible seaweeds. *Thyroid* 2004;14:836-41.

Correspondence Copyright © 2009 Massachusetts Medical Society.

#### INSTRUCTIONS FOR LETTERS TO THE EDITOR

Letters to the Editor are considered for publication, subject to editing and abridgment, provided they do not contain material that has been submitted or published elsewhere. Please note the following: • Letters in reference to a *Journal* article must not exceed 175 words (excluding references) and must be received within 3 weeks after publication of the article. Letters not related to a *Journal* article must not exceed 400 words. All letters must be submitted over the Internet at authors.NEJM.org. • A letter can have no more than five references and one figure or table. • A letter can be signed by no more than three authors. • Financial associations or other possible conflicts of interest must be disclosed. (Such disclosures will be published with the letters. For authors of *Journal* articles who are responding to letters, this information appears in the published articles.) • Include your full mailing address, telephone number, fax number, and e-mail address with your letter.

Our Web site: authors.NEJM.org

We cannot acknowledge receipt of your letter, but we will notify you when we have made a decision about publication. Letters that do not adhere to these instructions will not be considered. Rejected letters and figures will not be returned. We are unable to provide prepublication proofs. Submission of a letter constitutes permission for the Massachusetts Medical Society, its licensees, and its assignees to use it in the *Journal*'s various print and electronic publications and in collections, revisions, and any other form or medium.